


JC19 Rec'd PCT/PTO 23 MAY 2001

TRANSMITTAL LETTER TO THE UNITED STATES DESIGNATED/ELECTED OFFICE (DO/EO/US) CONCERNING A FILING UNDER 35 U.S.C. 371		ATTORNEY'S DOCKET NUMBER TEI-120
		U.S. APPLICATION NO. (known, see 37 CFR 1.101) <div style="font-size: 2em; font-weight: bold; text-align: center;">09/856556</div>
INTERNATIONAL APPLICATION NO. PCT/JP00/04146	INTERNATIONAL FILING DATE October 5, 2000	PRIORITY DATE CLAIMED October 5, 1999
TITLE OF INVENTION METHODS FOR INHIBITING OSTEOCLAST FORMATION OR METHODS FOR INHIBITING BONE RESORPTION		
APPLICANT(S) FOR DO/EO/US Yoshiaki AZUMA, Tomohiro OHTA, Keiji KOMORIYA, Akira KUDO, Sunao TAKSHITA, Keisuke KAJI		
<p>Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:</p> <ol style="list-style-type: none"> 1. <input checked="" type="checkbox"/> This is a FIRST submission of items concerning a filing under 35 U.S.C. 371. 2. <input type="checkbox"/> This is a SECOND or SUBSEQUENT submission of items concerning a filing under 35 U.S.C. 371. 3. <input checked="" type="checkbox"/> This express request to begin national examination procedures (35 U.S.C. 371(f) at any time rather than delay examination until the expiration of the applicable time limit set in 35 U.S.C. 371(b) and PCT Articles 22 and 39(I). 4. <input type="checkbox"/> A proper Demand for International Preliminary Examination was made by the 19th month from the earliest claimed priority date. 5. <input checked="" type="checkbox"/> A copy of the International Application as filed (35 U.S.C. 371(c)(2)) <ol style="list-style-type: none"> a. <input type="checkbox"/> is transmitted herewith (required only if not transmitted by the International Bureau). b. <input checked="" type="checkbox"/> has been transmitted by the International Bureau c. <input type="checkbox"/> is not required, as the application was filed in the United States Receiving Office (RO/US). 6. <input checked="" type="checkbox"/> A translation of the International Application into English (35 U.S.C. 371(c)(2)). 7. <input checked="" type="checkbox"/> Amendment to the claims of the International Application under PCT Article 19 (35 U.S.C. 371(c)(3)). <ol style="list-style-type: none"> a. <input type="checkbox"/> are transmitted herewith (required only if not transmitted by the International Bureau). b. <input type="checkbox"/> have been transmitted by the International Bureau. c. <input type="checkbox"/> have not been made; however, the time limit for making such amendment has NOT expired. d. <input checked="" type="checkbox"/> have not been made and will not be made. 8. <input type="checkbox"/> A translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371(c)(3)). 9. <input checked="" type="checkbox"/> An oath or declaration of the inventor(s) (35 U.S.C. 371(c)(4)). 10. <input type="checkbox"/> A translation of the annexes to the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371(c)(5)). <p>Items 11 to 16 below concern either document(s) or information included:</p> <ol style="list-style-type: none"> 11. <input checked="" type="checkbox"/> An Information Disclosure Statement under 37 CFR 1.97 and 1.98. 12. <input checked="" type="checkbox"/> An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included. 13. <input checked="" type="checkbox"/> A FIRST preliminary amendment. <input type="checkbox"/> A SECOND or SUBSEQUENT preliminary amendment. 14. <input type="checkbox"/> A substitute specification. 15. <input type="checkbox"/> A change of power of attorney and/or address letter. 16. <input type="checkbox"/> Other items or information. 		

3018 R&D PCT/PTO 2 3 MAY 2001

U.S. APPLICATION NO (If not, see 37 CFR 1.5) <div style="font-size: 2em; font-weight: bold; margin-left: 50px;">856556</div>		INTERNATIONAL APPLICATION NO <div style="font-weight: bold; margin-left: 50px;">PCT/JP00/04146</div>		ATTORNEY'S DOCKET NUMBER <div style="font-weight: bold; margin-left: 50px;">TEI-120</div>	
17. <input checked="" type="checkbox"/> The following fees are submitted: <div style="margin-left: 20px;"> Basic National Fee (37 CFR 1.49(a)(1)-(5)): Search Report has been prepared by the EPO or JPO..... International preliminary examination fee paid to USPTO (37 CFR 1.482) No international preliminary examination fee paid to USPTO (37 CFR 1.482) but international search fee paid to USPTO (37 CFR 1.445(a)(2))..... Neither international preliminary examination fee (37 CFR 1.482) nor international search fee (37 CFR 1.445(a)(2)) paid to USPTO..... International preliminary examination fee paid to USPTO (37 CFR 1.482) and all claims satisfied provisions of PCT Article 33(2)-(4)..... </div>				CALCULATIONS	PTO USE ONLY
<div style="text-align: right; font-weight: bold;">ENTER APPROPRIATE BASIC FEE AMOUNT =</div>				<div style="font-weight: bold;">\$ 860.00</div>	
Surcharge of \$130.00 for furnishing the oath or declaration later than <input type="checkbox"/> 20 <input type="checkbox"/> 30 months from the earliest claimed priority date (37 CFR 1.49(e)).				\$	
Claims	Number Filled	Number Extra	Rate		
Total Claims	6-20 =	0	X \$18	\$ 0	
Independent Claims	2-3 =	0	X \$78	\$ 0	
Multiple dependent claim(s) (if applicable)			+ \$260	\$ 0	
TOTAL OF ABOVE CALCULATIONS				=	\$860.00
Reduction by 1/2 for filing by small entity, if applicable. Verified Small Entity statement must also be filed. (Note 37 CFR 1.9, 1.27, 1.28)				\$	
SUBTOTAL				=	\$860.00
Processing fee of \$130.00 for furnishing the English translation later than <input type="checkbox"/> 20 <input type="checkbox"/> 30 months from the earliest claimed priority date (37 CFR 1.49(f)).				+	\$
TOTAL NATIONAL FEE				=	\$860.00
Fee for recording the enclosed assignment (37 CFR 1.21(h)). The assignment must be accompanied by an appropriate sheet (37 CFR 3.28, 3.31). \$40.00 per property				+	\$ 0
TOTAL FEES ENCLOSED				=	\$860.00
				Amount to be refunded	\$
				Charged	\$860.00
a. <input type="checkbox"/> A check in the amount of \$_____ to cover the above fees is enclosed. b. <input checked="" type="checkbox"/> Please charge my Deposit Account No <u>18-0013</u> in the amount of <u>\$860.00</u> to cover the above fees. A duplicate of this sheet is enclosed. c. <input checked="" type="checkbox"/> The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any overpayment to Deposit Account No. <u>18-0013</u> . A duplicate copy of this sheet is enclosed.					
NOTE: Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (37 CFR 1.137(a) or (b)) must be filed and granted to restore the application to pending status.					
SEND ALL CORRESPONDENCE TO: <div style="display: flex; justify-content: space-between;"> <div style="width: 60%;"> David K. Benson Rader, Fishman & Grauer, L.P.C. 1233 20th Street, N.W. Suite 501 Washington, DC 20036 202-955-3750 </div> <div style="width: 35%; text-align: right;"> <div style="font-size: 1.5em; margin-bottom: 5px;">  </div> <div style="font-weight: bold; margin-bottom: 10px;">DAVID K. BENSON</div> <div style="font-weight: bold;">REGISTRATION NUMBER 42,314</div> </div> </div>					
Dated: May 23, 2001					

TEI-118

09/856556

PATENT APPLICATION

IC18 Rec'd PCT/PTO 23 MAY 2001

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In the U.S. Nationalization

Application of PCT/JP00/04146

Yoshiaki AZUMA et al.

Attn: Application Branch

Application No. To Be Assigned

Filed: May 22, 2001

For: METHODS FOR INHIBITING OSTEOCLAST FORMATION OR METHODS FOR
INHIBITING BONE RESORPTION

PRELIMINARY AMENDMENT

Commissioner of Patents
Washington, DC 20231

Sir:

Prior to the initial examination, please amend the above-identified application as follows:

IN THE CLAIMS:

Please amend claim 5 and add new claims 7 and 8 as follows:

5. (Amended) The method for inhibiting bone resorption as described in claim 3 wherein said co-culture comprises an inducing factor of osteoclast formation and/or bone resorption in the co-culture medium.

7. The method for inhibiting bone resorption as described in claim 4 wherein said co-culture comprises an inducing factor of osteoclast formation and/or bone resorption in the co-culture medium.

8. The method for inhibiting osteoclast formation or the method for inhibiting bone resorption according to claim 7 wherein said inducing factor comprises at least one factor


TEI-118-09856556

selected from the group consisting of macrophage colony-stimulating factor, osteoclast forming factor, tumor necrosis factor, interleukin-1, interleukin-3, interleukin-6, interleukin-11, interleukin-15, interleukin-17, prostaglandins, parathyroid hormone, parathyroid hormone-related peptide, granulocyte macrophage colony-stimulating factor, active vitamin D and its derivatives.

REMARKS

This preliminary amendment is requested to change all multiple dependency claims to single dependency claims to reduce the initial filing fee. No new matter has been added. Entry of this amendment is requested.

Respectfully submitted,



David K. Benson
Reg. No. 42,314

Date: May 23, 2001

RADER, FISHMAN & GRAUER, PLLC

The Lion Building
1233 20th Street, N.W., Suite 501
Washington, D.C. 20036
Tel: (202) 955-3750
Fax: (202) 955-3751
Customer No. 23353

MARKED-UP VERSION TO SHOW CHANGES MADE

IN THE CLAIMS:

Please amend claim 5 as follows:

5. (Amended) The method for inhibiting bone resorption as described in claim 3 [or 4] wherein said co-culture comprises an inducing factor of osteoclast formation and/or bone resorption in the co-culture medium.

09/856556

4/PRTS

09/856556 T-412

JC18 Rec'd PCT/PTO 23 MAY 2001

- 1 -

DESCRIPTION

METHOD FOR INHIBITING OSTEOCLAST FORMATION OR METHOD FOR INHIBITING BONE RESORPTION

5

Technical Field

The present invention relates to a method for inhibiting osteoclast formation, or a method for inhibiting bone resorption.

10 More particularly this invention relates to a method for inhibiting osteoclast formation or a method for inhibiting bone resorption by ultrasound (which may be called as "US" hereinafter) treatment.

Background Art

15 A known and conventional method for inhibiting osteoclast formation or for inhibiting bone resorption includes a method consisting of adding a bisphosphonate to a co-culture system [Bilezikian, JP., Raisz, LG., and Rodan, GA. eds., "Principles of bone biology," Academic Press, New York (1996)]. Clinically, a medicine effective in inhibiting osteoclast formation or inhibiting bone resorption has been used for treating of a bone disease caused by
20 abnormal bone resorption due to osteoclast [Bilezikian, JP., Raisz, LG., and Rodan, GA. eds., "Principles of bone biology," Academic Press, New York (1996)].

In the method for adding or administering of a medicine or pharmaceuticals, however, the medicine or pharmaceuticals are taken into
25 body or cells directly, and thus it is not free from the risk of causing some side-effects. Therefore, there is a demand for a method whereby it is possible to inhibit osteoclast formation or bone resorption, while being less likely to cause side-effect than does the conventional method.

30 In view of this, the present inventors had tried hard to find such a method as mentioned above, that is, a method whereby it is possible to inhibit osteoclast formation or bone resorption, in spite of being less likely to cause side-effect than does the conventional method, and had observed that stimulation with ultrasound, which is physical and non invasive, can effectively inhibit osteoclast formation or bone resorption, and achieved this

09856556-05201

invention.

Disclosure of the Invention

The present invention provides a method for inhibiting osteoclast
5 formation or a method for inhibiting bone resorption which comprises
exposing cells to ultrasound in the culture containing an osteoclast precursor
and an inducing factor of osteoclast formation and /or bone resorption, or a
method for inhibiting osteoclast formation or a method for inhibiting bone
resorption whici comprises exposing cells to ultrasound in the co-culture
10 containing an osteoclasst precursor and a supporting cell for
osteoclastogenesis.

Brief Description of the Drawings

Figure 1 shows the number of osteoclasts over time in an experiment
15 where the development of mouse osteoclast precursors into osteoclasts is
stimulated by sRANKL.

Figure 2 shows the number of osteoclasts obtained when mouse
osteoclast precursors have been co-cultivated with mouse bone marrow cells
for 96 hours.

Figure 3 shows the number of osteoclasts obtained when mouse
20 osteoclast precursors have been co-cultivated with mouse bone marrow cells
for 120 hours.

Figure 4 shows the area of resorption pits by osteoclasts which have
been derived from mouse oseteoclast precursors co-cultivated with mouse
25 bone marrow cells.

Note that, for all the relevant Figures, * represents the test result is
significantly different from the result of control group at $p < 0.05$ based on the
Student's t-test, while *** represents the same at $p < 0.001$.

30 Best Mode for Carrying Out the Invention

The present invention is described below in detail.

The osteoclast precursor used in this invention is the precursor that
develop into osteoclasts in the presence of an inducing factor added to the
culture medium, or in the presence of supporting cells added to the culture

medium which have been known from the effect to other osteoclast precursors to have an activity to support osteoclast formation. As such osteoclast precursors, bone marrow-derived precursors, spleen-derived precursors and blood monocyte-derived precursors may be mentioned.

5 Those precursor cells may be obtained from the bone marrow, spleen or peripheral blood of humans, mice, etc., and they can be cultivated through passage on a culture medium specially prepared for the emphasized growth of such osteoclast precursors.

The inducing factor which stimulates osteoclast precursors to turn them
10 into osteoclasts is at least one chosen from the group comprising a macrophage colony stimulating factor (to be referred to as "M-CSF" hereinafter), osteoclast forming factor (RANKL/TRANSE/ODF/OPGL, or to be referred to as "RANKL" hereinafter), tumor necrosis factor (the same as above "TNF"), interleukin 4 (IL-4), and vascular endothelial cell growth
15 factor (VEGF). Those factors may be used alone or in combination. RANKL is preferably used, or an artificially synthesized soluble RANKL is most preferred when used at a concentration of 1 to 250 ng/ml. Addition of the inducing factor into the culture medium should be occurred during a period immediately after the inoculation of osteoclast precursors onto the medium
20 till osteoclast formation .

Besides the osteoclast precursors, the co-culture system may include at least one chosen from the group comprising osteoblasts, stroma cells, fibroblasts, T-cells and B-cells. A single kind or several kinds of those cells may be included.

25 Moreover, the co-culture system preferably includes an inducing factor. The inducing factor may include, in addition to M-CSF, RANKL, TNF, IL-4 and VEGF mentioned above, interleukin-1 (to be referred to as "IL-1" hereinafter), interleukin-3 (the same as above "IL-3"), interleukin-6 (the same as above "IL-6"), interleukin-11 (the same as above "IL-11"),
30 interleukin-15 (the same as above "IL-15"), interleukin-17 (the same as above "IL-17"), prostaglandins (the same as above "PG") including prostaglandin E2, parathyroid hormone (the same as above "PTH"), parathyroid hormone-related peptide (the same as above "PTHrP"), granulocyte/macrophage colony stimulating factor (the same as above

"GM-CSF"), and activated vitamin Ds including $1\alpha,25$ -dihydroxyvitamin D3 (the same as above "VD"). They may be used alone or in combination.

Of all the inducing factors cited above, RANKL is preferred to be used. This is because RANKL is a factor essential for the osteoclast formation, and
5 plays an important role in bone metabolism/maintenance [Takahashi, N., et al., Biochem Biophys Res Comm 256: 449-455 (1999)]. RANKL is mainly present on the surface of cells known to support the osteoclast formation, such as osteoblasts, stroma cells, fibroblasts, etc., and induces osteoclast formation by binding with receptors (RANK/TRANR) of the osteoclast
10 precursors [Takahashi, N., et al., Biochem Biophys Res Comm 256: 449-455 (1999)]. Because RANKL hardly exist as a soluble factor in the body, the co-culture system for the osteoclast formation may be thought as a system resembling an in-situ environment involved in the bone metabolism. RANKL plays an important role in the metabolism of a normal bone, but it is
15 also deeply involved in the development of bone-related diseases. For example, in postmenopausal osteoporosis, the number of osteoclasts and their bone resorbing activity are increased, and the bone mass decreased. It is thought that RANKL is involved in the growth of osteoblasts [Kong, YY. et al., Nature 397:315-323 (1999)]. It has been demonstrated that a biological
20 molecule called osteoclast formation inhibiting factor (to be referred to as "OPG/OCIF" herinafter) that interferes with the mutual binding of RANKL molecules not only promotes the osteoclast formation stimulated by RANKL, but also, in animal models having the diseases resulting from the enhanced osteoclast formation or enhanced activity of osteoclasts with bone-related
25 diseases such as hypercalcemia, osteoporosis, etc., inhibits the osteoclast formation, and also improves the symptoms such as bone mass reduction and increase of calcium level in blood, by suppressing the bone absorbing activity. [Lacey DL., et al., Cell 93: 165-176 (1998)]. If it becomes possible to develop a method by which to inhibit the osteoclast formation stimulated by
30 RANKL or induced via a certain agent stimulated by RANKL, as a therapeutic means to recover the reduced bone mass or bone disruption in bone-related diseases resulting from the enhanced bone resorption such as osteoporosis, chronic rheumatoid arthritis, etc.; the reduced bone mass or the loosening after arthroplasty; or the disruption of a bone as a result of cancer

metastasized to the bone, etc., the method will be prospective as a therapy to treat bone-related diseases.

The co-culture system used in this invention is a culture system in which osteoclast precursors and at least one type of cells which support osteoclast formation are cultivated in the same medium. The cell to support
5 osteoclast formation is at least one cell selected from the group consisting of osteoblasts, stroma cells, fibroblasts, T-cells and B-cells. They may be used alone or in combination. Of those cells, stroma cells are preferably used, and bone marrow-derived stroma cells are more preferred. The most
10 preferred combination consists of bone-marrow derived osteoclast precursors and bone marrow-derived stroma cells. The bone marrow-derived stroma cells may include mouse bone marrow-derived stroma cells such as ST-2, TSB-13-9, etc.

The cell line ST-2 is available from American Type Culture Collection,
15 and it can be cultivated and maintained on the culture medium that will be detailed later, to be ready for experimentation.

The medium used in this invention to cultivate osteoclast precursors may include alpha MEM medium containing 8 to 1000 ng/ml of M-CSF and supplemented with 5 - 15% fetal bovine serum. Particularly, the alpha
20 MEM medium containing 100 ng/ml of M-CSF and supplemented with 10% fetal bovine serum is preferred.

The medium used for the co-culture system may include alpha MEM medium supplemented with an inducing factor described above, and 5 - 15% fetal bovine serum. Particularly, alpha MEM medium supplemented with
25 $1 \alpha, 25$ -dihydroxy vitamin D3 having a concentration of 1×10^{-10} M to 1×10^{-7} M, and 10% fetal bovine serum is preferred. The inducing factor may be added to the medium during a period between a time when osteoclast precursors and osteoclast formation supporting cells such as bone marrow-derived stroma cells are inoculated concomitantly into the medium
30 and a time when osteoclasts are formed.

The ultrasound to be used in this invention includes ultrasound with an output too low to generate heat. More specifically, the ultrasound is a low-output pulsed ultrasound consisting of a series of bursts of which the wave component in a burst has a frequency of 1.3 to 2 MHz; the repetition

frequency of bursts is 100 to 1000 KHz; the width of a burst is 10 to 2000 μ sec; and the output intensity is 100 mW/cm² or less. Exposure of the ultrasound is achieved, to cite an example for illustration, as follows: an ultrasound generator provided by Exogen is so adjusted as to give low-output
5 ultrasound pulses lasting 20 minutes in total for a day; the generator is so positioned with respect to the culture as to expose the wave to the bottom of the culture well; and the exposure continues for three to five days. The preferred properties of the ultrasound exposure, for example, are as follows; the width of a burst was 200 μ sec, the frequency was 1.5 MHz, the repetition
10 frequency of bursts was 1 kHz and the output intensity was 30 mW/cm².

The ultrasound generator to be used in this invention is not limited to any specific one, but, if clinically used, what is preferred is, for example, a generator enabling basic, non-invasive therapy as disclosed in US patent (No. 4,530,360) conferred to Duarte, which is capable of applying pulsed
15 ultrasound percutaneously from a point close to the site to be treated. The preferred ultrasound emitted by such a generator may consist of a series of bursts of which the wave component in a burst has a frequency of 1.3 to 2 MHz; the repetition frequency of bursts is 100 to 1,000 Hz; the width of a burst is 10 to 2,000 μ sec; and the output intensity is 100 mW/cm² or less.
20 The duration of exposure is preferably shorter than 20 minutes for a day. Or, an ultrasound-based therapeutic system incorporating a body applicator unit communicable with a remote control unit as disclosed in US Patent (No. 5,003,965) conferred to Talish et al. may be used. The ultrasound in this case is a low-output pulsed ultrasound with a frequency of 1.3 to 2 MHz and
25 an output intensity of 1 to 50 mW/cm².

According to the method of this invention for inhibiting the osteoclast formation or bone resorption, it will be possible to expose ultrasound onto a lesion where bone mass reduction or bone disruption occurs as a result of the enhanced osteoclast formation or enhanced bone resorption, and, if it is
30 clinically applied, to recover the reduced bone mass or bone disruption in bone-related diseases such as osteoporosis, chronic rheumatoid arthritis, etc., to fix aseptic loosening after arthroplasty, and to treat the bone disruption due to a tumor metastasized to a bone. Moreover, the method of this invention for the inhibiting osteoclast formation or bone resorption is

Examples

Examples 1-1 to 1-4, and Comparative Examples 1-1 to 1-4

Comparative Example 1-1: control group cultivated for 72 hours.

20 Example 1-1: ultrasound exposure group cultivated for 72 hours.

For this group, ultrasound was exposed from the bottom of each well for 20 minutes daily for two days successively.

Comparative Example 1-1: control group cultivated for 72 hours.

20 **Example 1-1:** ultrasound exposure group cultivated for 72 hours.
For this group, ultrasound was exposed from the bottom of each well for 20
minutes daily for two days successively.

Comparative Example 1-2: control group cultivated for 96 hours.

Example 1-2: ultrasound exposure group cultivated for 96 hours.
 25 For this group, ultrasound was exposed from the bottom of each well for 20
 minutes daily for three days successively.

Comparative Example 1-3: control group cultivated for 108 hours.

Example 1-3: ultrasound exposure group cultivated for 108 hours. For this group, ultrasound was exposed from the bottom of each well for 20 minutes daily for four days successively.

Comparative Example 1-4: control group cultivated for 144 hours.

Example 1-4: ultrasound exposure group cultivated for 144 hours. For this group, ultrasound was exposed from the bottom of each well for 20 minutes daily for five days successively.

For both the Comparative Examples and Examples, sRANKL was dissolved in sterilized distilled water to give a concentration of 10 µg/ml; and the resulting solution was added so as to be diluted 200 times to give a concentration of 50 ng/ml. On the third day, the media of both the Comparative Examples and Examples were replaced with fresh 10% FBS-aMEM containing 50 ng/ml of sRANKL.

Exposure of ultrasound was not applied to the Comparative Examples 1-1 to 1-4. For Examples 1-1 to 1-3, a low output ultrasound was applied from the bottom of each well for 20 minutes daily using an ultrasound generating unit provided by Exogen which had been modified to be adapted for cell culture. Example 1-1 received ultrasound exposure for two days; Example 1-2 for three days; Example 1-3 for four days and Example 1-4 for five days successively. As to the output properties of the ultrasound applied, the width of a burst was 200 µsec, the frequency was 1.5 MHz, repetition frequency was 1 kHz and the output was 30 mW/cm².

Removal of the culture medium took place 72 hours after the onset of cultivation for Comparative Example 1-1 and Example 1-1, 96 hours after for Comparative Example 1-2 and Example 1-2, 108 hours after for Comparative Example 1-3 and Example 1-3, and 144 hours after for Comparative Example 1-4 and Example 1-4. The cells were fixed with phosphate-buffered formalin, stained with tartaric acid-resistant acidic phosphatase (TRAP) according to convention for identification of osteoclasts, and placed under an optical microscope, and the number of osteoclasts was determined on the premise that a TRAP-positive polynucleated cell having ten or more nuclei is an osteoclast.

The results are shown in Figure 1. As is obvious from the Figure, the numbers of osteoclasts in Examples 1-2 to 1-4 are less than those of Comparative Examples 1-2 to 1-4.

Examples 2-1 to 2-2 and Comparative Examples 2-1 to 2-2

MDBMs were obtained by extracting the bone marrow of femur and tibia of mice, separating monocytes therefrom according to convention, inoculating the monocytes on dishes to give a concentration of 4×10^6 cells/10 cm-dish, and selectively cultivating adhesive cells in the presence of 100

ng/ml of M-CSF for three days. The culture medium was an alpha MEM medium containing 100 ng/ml of M-CSF and supplemented with 10% fetal bovine serum. The cell line TSB-13-9 derived from mouse bone marrow-derived stroma cells is an established cell line prepared from cells
5 obtained from the bone marrow of SV-40-large-T-antigen transgenic mice. The cells of TSB-13-9 were inoculated on petri dishes to give a concentration of 2×10^5 cells/10 cm-dish, and cultivated for three days. The medium was an alpha MEM medium supplemented with 10% fetal bovine serum.

Cells of MDBM and TSB-13-19 were simultaneously inoculated into a
10 6-well plate 2 ml for each well to give concentrations of 1×10^5 and 1×10^6 cells/well, respectively. To each well, 1×10^{-8} M $1\alpha,25$ -dihydroxyvitamin D3 was added. For the experiment, following groups were prepared.

Comparative Example 2-1: control group cultivated for 96 hours.

Example 2-1: ultrasound exposure group cultivated for 96 hours.
15 For this group, ultrasound was exposed from the bottom of each well for 20 minutes daily for three days successively.

Comparative Example 2-2: control group cultivated for 120 hours.

Example 2-2: ultrasound exposure group cultivated for 120 hours.
For this group, ultrasound was exposed from the bottom of each well for 20
20 minutes daily for four days successively.

For both the Comparative Examples and Examples, $1\alpha,25$ -dihydroxyvitamin D3 was dissolved in ethanol to give a concentration of 1×10^{-5} M; and the resulting solution was added so as to be diluted 1,000 times to give a concentration of 1×10^{-8} M. On the third day, the media of
25 both the Comparative Examples and Examples were replaced with fresh 10% FBS - α MEM containing 1×10^{-8} M $1\alpha,25$ -dihydroxyvitamin D3.

Exposure of ultrasound was not applied to Comparative Examples 1-1 and 1-2. For Examples 1-1 and 1-2, low output ultrasound was applied from the bottom of each well for 20 minutes daily using an ultrasound generating
30 unit provided by Exogen which had been modified to be adapted for cell culture. Example 1-1 received the ultrasound exposure for three days successively, and Example 1-2 for four days. The ultrasound consisted of the output properties, the width of a burst was 200 μ sec, the frequency was 1.5 MHz, the repetition frequency of bursts was 1 kHz and the output

intensity was 30 mW/cm².

Removal of the medium took place 96 hours after the onset of cultivation for Comparative Example 1-1 and Example 1-1, and 120 hours for Comparative Example 1-2 and Example 1-2. The cells were fixed with 10% phosphate-buffered formalin, stained with tartaric acid resistive, acid phosphatase (TRAP) according to convention for identification of osteoclasts, and placed under an optical microscope, and the number of osteoclasts was determined on the premise that a TRAP-positive polynucleated cell having ten or more nuclei is an osteoclast. The results are shown in Figure 1.

As is obvious from Figures 2 and 3, the numbers of osteoclasts in Examples 1-1 and 1-2 are less than those of Comparative Examples 1-1 and 1-2.

Examples 3-1 and Comparative Example 3-1

MDBMs prepared as in Example 1 were inoculated onto a 24-well plate 1 ml for each well to give a concentration of 2×10^4 cells/well. To each well of the 24-well plate, a hydroxyapatite-coated disc (Osteologic, Millennium Biologix, Inc.) had been placed in advance to receive the cells to grow them thereupon. sRANKL was added to each well to give a concentration of 50 ng/ml. sRANKL had been obtained from Pepro-Tech.

For the experiment, following groups were prepared.

Comparative Example 3-1: control group cultivated for 120 hours.

Example 3-1: ultrasound exposure group cultivated for 120 hours. For this group, ultrasound was exposed from the bottom of each well for 20 minutes daily for four days successively.

For both the Comparative Examples and Examples, sRANKL was dissolved in sterilized distilled water to give a concentration of 10 µg/ml; and the resulting solution was added so as to be diluted 200 times to give a concentration of 50 ng/ml. On the third day, the media of both the Comparative Examples and Examples were replaced with fresh 10% FBS - α MEM containing 50 ng/ml of sRANKL.

Exposure of ultrasound was not applied to Comparative Example 3-1. For Example 3-1, a low output ultrasound was applied from the bottom of each well for 20 minutes daily for four days successively using a ultrasound generating unit provided by Exogen which had been modified to be adapted

for cell culture. The ultrasound consisted of the output properties, the width of a burst was 200 μ sec, the frequency was 1.5 MHz, the repetition frequency of bursts was 1 kHz and the output intensity was 30 mW/cm².

Removal of the medium took place 120 hours after the onset of cultivation for both Comparative Example 3-1 and Example 3-1, and the hydroxyapatite-coated discs were removed. The disc was washed with distilled water, and placed under an optical microscope, and absorption of the hydroxyapatite mass by osteoclasts was checked. A picture of a part of the disc (whose area is 50% or more of the total area) was fed through an image processor into a computer, and the area of disc undergoing resorption was determined by means of an image analysis application. The resorbed area relative to the total was determined and the results are shown in Figure 3.

As is obvious from Figure. 3, the relative area of disc subjected to osteoclastic resorption is smaller in Example 3-1 than in Comparative Example 3-1.

Example 4-1 and Comparative Example 4-1

MDBM and the cell line TSB-13-9 derived from mouse bone-marrow derived stroma cells were prepared as in Example 2. The medium was an alpha MEM medium supplemented with 10% fetal bovine serum .

MDBM and TSB-13-19 were inoculated at the same time into a 24-well plate to give concentrations of 2×10^4 and 1×10^5 cells/well, respectively. The volume the culture medium was arranged as 1 ml finally. To each well of the 24-well plate, a hydroxyapatite-coated disc (Osteologic, Millennium Biologix, Inc.) had been placed in advance to receive the cells to grow them thereupon. $1 \alpha, 25$ -dihydroxyvitamin D₃ was added to each well to give a concentration of 1×10^{-8} M.

For the experiment, following groups were prepared.

Comparative Example 4-1: control group cultivated for 120 hours.

Example 4-1: ultrasound exposure group cultivated for 120 hours. For this group, ultrasound was exposed from the bottom of each well for 20 minutes daily for four days successively.

For both the Comparative Example and Example, $1\alpha, 25$ -dihydroxyvitamin D₃ was dissolved in ethanol to give a concentration

of $1 \times 10^{-5} \text{M}$; and the resulting solution was added so as to be diluted 1,000 times to give a concentration of $1 \times 10^{-8} \text{M}$. On the third day, the media of both the Comparative Example and Example were replaced with fresh 10% FBS- α MEM containing $1 \times 10^{-8} \text{M}$ $1\alpha,25$ -dihydroxyvitamin D3.

5 Exposure of ultrasound was not applied to Comparative Example 4-1. For Example 4-1, low output ultrasound was applied from the bottom of each well for 20 minutes daily for four days successively using an ultrasound generating unit provided by Exogen which had been modified to be adapted for cell culture. The ultrasound consisted of the output properties, the width
10 of a burst was 200 μsec , the frequency was 1.5 MHz, the repetition frequency of bursts was 1 kHz and the output intensity was 30 mW/cm².

Removal of the medium took place 120 hours after the onset of cultivation for both Comparative Example 4-1 and Example 4-1. The disc was washed with distilled water, and placed under an optical microscope,
15 and resorption of the hydroxyapatite mass by osteoclasts was checked. A picture of a part of the disc (whose area is 50% or more of the total area) was fed through an image processor into a computer, and the area of disc undergoing absorption was determined by means of an image analysis application. The resorbing area relative to the total was determined and the
20 results are shown in Figure 4.

As is obvious from Figure 4, the relative area of disc subjected to osteoclastic resorption is smaller in Example 4-1 than in Comparative Example 4-1.

The present invention provides a method for inhibiting osteoclast
25 formation or a method for inhibiting bone resorption, characterized in using the exposure of ultrasound. A clinical application to the medical treatment for bone-related deceases, such as a bone mass reduction or a bone disruption by osteoporosis and chronic rheumatoid arthritis, a bone destruction of aseptic loosening with joint arthroplasty just after the replacement with an
30 artificial joint, by ultrasound exposure, which is physical and non-invasive, to a lesion where bone mass reduction or bone destruction occurs as a result of the enhanced osteoclast formation or enhanced bone resorption, without some side-effects, becomes possible. Moreover, the clinical application of the method of this invention for inhibiting the osteoclast formation or bone

[illegible]

CLAIMS

1. A method for inhibiting osteoclast formation or a method for inhibiting bone resorption which comprises exposing cells to ultrasound in the culture
5 containing an osteoclast precursor and an inducing factor of osteoclast formation and /or bone resorption.
2. The method for inhibiting osteoclast formation or the method for inhibiting bone resorption according to claim 1 wherein said inducing factor comprises at least one factor selected from the group consisting of
10 macrophage colony-stimulating factor, osteoclast differentiating factor, tumor necrosis factor, interleukin-4, and vascular endothelial cell growth factor.
3. A method for inhibiting osteoclast formation or a method for inhibiting bone resorption whici comprises exposing cells to ultrasound in the co-culture containing an osteoclast precursor and an supporting cell for osteogenesis.
- 15 4. The method for inhibiting osteoclast formation or the method for inhibiting bone resorption according to claim 3 wherein said co-culture system comprises at least one cell seledcted from the group consisting of osteoblast, stroma cell, fibroblast, T-lymphocyte and B-lymphocyte.
5. The method for inhibiting osteoclast formation or the method for
20 inhibiting bone resorption as described in claim 3 or 4 wherein said co-culture comprises an inducing factor of osteoclast formation and/or bone resorption in the co-culture medium.
6. The method for inhibiting osteoclast formation or the method for inhibiting bone resorption according to claim 5 wherein said inducing factor
25 comprises at least one factor selected from the group consisting of macrophage colony-stimulating factor, osteoclast forming factor, tumor necrosis factor, interleukin-1, interleukin-3, interleukin-6, interleukin-11, interleukin-15, interleukin-17, prostaglandins, parathyroid hormone, parathyroid hormone-related peptide, granulocyte macrophage colony
30 -stimulating factor, active vitamin D and its derivatives.

ABSTRACT

A method for inhibiting osteoclast formation or a method for inhibiting bone resorption which comprises exposing cells to ultrasound in the culture containing an osteoclast precursor and an inducing factor of osteoclast formation and /or bone resorption. Or, a method for inhibiting osteoclast formation or a method for inhibiting bone resorption which comprises exposing cells to ultrasound in the co-culture containing an osteoclast precursor and a supporting cell for osteoclastogenesis.

10

Figure 1




Figure 2




Figure 3





Figure 4



Declaration and Power of Attorney For Patent Application

特許出願宣言書及び委任状

Japanese Language Declaration

日本語宣言書

下記の氏名の発明者として、私は以下の通り宣言します。 As a below named inventor, I hereby declare that:

私の住所、私書箱、国籍は下記の私の氏名の後に記載された通りです。

My residence, post office address and citizenship are as stated next to my name.

下記の名称の発明に関して請求範囲に記載され、特許出願している発明内容について、私が最初かつ唯一の発明者（下記の氏名が一つの場合）もしくは最初かつ共同発明者であると（下記の名称が複数の場合）信じています。

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled

"METHODS FOR INHIBITING OSTEOCLAST FORMATION"

OR METHODS FOR INHIBITING BONE RESORPTION"

上記発明の明細書（下記の欄で x 印がついていない場合は、本書に添付）は、 the specification of which is attached hereto unless the following box is checked:

☐ __月__日に提出され、米国出願番号または特許協定条約
国際出願番号を____とし、
(該当する場合) _____ に訂正されました。
☒ was filed on June 23, 2000
as United States Application Number or
PCT International Application Number
PCT/JP00/04146 and was amended on
_____ (if applicable).

私は、特許請求範囲を含む上記訂正後の明細書を検討し、内容を理解していることをここに表明します。

I hereby state that I have reviewed and understand the contents of the above identified specification, including the claims, as amended by any amendment referred to above.

私は、連邦規則法典第 37 編第 1 条 56 項に定義されるとおり、特許資格の有無について重要な情報を開示する義務があることを認めます。

I acknowledge the duty to disclose information which is material to patentability as defined in Title 37, Code of Federal Regulations, Section 1.56.

Japanese Language Declaration (日本語宣言書)

私は、米国法典第35編119条(a)-(d)項又は365条(b)項に基づき下記の、米国以外の国の少なくとも一カ国を指定している特許協力条約365(a)項に基づく国際出願、又は外国での特許出願もしくは発明者証の出願について外国優先権をここに主張するとともに、優先権を主張している、本出願の前に出願された特許または発明者証の外国出願を以下に、枠内をマークすることで、示しています。

I hereby claim foreign priority under Title 35, United States Code, Section 119(a)-(d) or 365(b) of any foreign application(s) for patent or inventor's certificate, or 365(a) of any PCT International application which designated at least one country other than the United States, listed below and have also identified below, by checking the box, any foreign application for patent or inventor's certificate, or PCT International application having a filing date before that of the application on which priority is claimed.

Prior Foreign Application(s)

外国での先行出願

Priority Claimed

優先権主張

11-284075

Japan

5/October/1999

(Number)
(番号)(Country)
(国名)(Day/Month/Year Filed)
(出願年月日)
☒ ☐
Yes No
はい いいえ
(Number)
(番号)(Country)
(国名)(Day/Month/Year Filed)
(出願年月日)
☐ ☐
Yes No
はい いいえ

私は、第35編米国法典第119条(e)項に基づいて下記の米国特許出願規定に記載された権利をここに主張いたします。

I hereby claim the benefit under Title 35, United States Code, Section 119(e) of any United States provisional application(s) listed below.

(Application No.)
(出願番号)(Filing Date)
(出願日)(Application No.)
(出願番号)(Filing Date)
(出願日)

私は、下記の米国法典第35編120条に基づいて下記の米国特許出願に記載された権利、又は米国を指定している特許協力条約365条(c)に基づく権利をここに主張します。また、本出願の各請求範囲の内容が米国法典第35編112条第1項又は特許協力条約で規定された方法で先行する米国特許出願に開示されていない限り、その先行米国出願書提出日以降で本出願書の日本国内または特許協力条約国際提出日までの期間中に入手された、連邦規則法典第37編1条56項で定義された特許資格の有無に関する重要な情報について開示義務があることを認識しています。

I hereby claim the benefit under Title 35, United States Code, Section 120 of any United States application(s), or 365(c) of any PCT International application designating the United States, listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States or PCT International application in the manner provided by the first paragraph of Title 35, United States Code Section 112, I acknowledge the duty to disclose information which is material to patentability as defined in Title 37, Code of Federal Regulations, Section 1.56 which became available between the filing date of the prior application and the national or PCT International filing date of application.

(Application No.)
(出願番号)(Filing Date)
(出願日)(Status: Patented, Pending, Abandoned)
(現況: 特許許可済、係属中、放棄済)(Application No.)
(出願番号)(Filing Date)
(出願日)(Status: Patented, Pending, Abandoned)
(現況: 特許許可済、係属中、放棄済)

私は、私自身の知識に基づいて本宣言書中で私が行なう表明が真実であり、かつ私の入手した情報と私の信じているところに基づく表明が全て真実であると信じていること、さらに故意になされた虚偽の表明及びそれと同等の行為は米国法典第18編第1001条に基づき、罰金または拘禁、もしくはその両方により処罰されること、そしてそのような故意による虚偽の声明を行なえば、出願した、又は既に許可された特許の有効性が失われることを認識し、よってここに上記のごとく宣言を致します。

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

Japanese Language Declaration (日本語宣言書)

委任状: 私は下記の発明者として、本出願に関する一切の
手続きを米特許商標局に対して遂行する弁理士または代理人
として、下記の者を指名いたします。(弁護士、または代理
人の氏名及び登録番号を明記のこと)

John E. McGarry	22,360
H. Lawrence Smith	24,900
Ralph T. Rader	28,772
Joseph V. Coppola, Sr.	33,373
Michael B. Stewart	36,018
Alexander D. Rabinovich	37,425
Kevin D. Rutherford	40,412
Glenn E. Forbis	40,610
Ronald P. Kananen	24,104
Matthew J. Russo	41,282
G. Thomas Williams	42,228

POWER OF ATTORNEY: As a named inventor, I hereby appoint
the following attorney(s) and/or agent(s) to prosecute this
application and transact all business in the Patent and Trademark
Office connected therewith (list name and registration number).

Joel E. Bair	33,356
Richard D. Grauer	22,388
Michael D. Fishman	31,951
Mark A. Davis	37,118
Stefan V. Chmielewski	39,914
Shumel Livnat	33,949
Kristin L. Murphy	41,212
David K. Benson	42,314
Christopher M. Tanner	41,518
Robert S. Green	41,800

書類送付先

RADER, FISHMAN & GRAUER PLLC
1233 20th Street, N.W.
Suite 501
Washington, D.C. 20036

Send Correspondence to:

RADER, FISHMAN & GRAUER PLLC
1233 20th Street, N.W.
Suite 501
Washington, D.C. 20036

直接電話連絡先: (名前及び電話番号)

(202) 955-3750

Direct Telephone Calls to: (name and telephone number)

(202) 955-3750

唯一または第一発明者名	1 - 00	Full name of sole or first inventor	Yoshiaki AZUMA
発明者の署名	日付	Inventor's signature	Date May 14, 2001
住所		Residence	TOKYO, JAPAN JPX
国籍		Citizenship	Japanese
私書箱		Post Office Address	c/o Teijin Limited, Tokyo Research Center, 3-2, Asahigaoka 4-chome, Hino-shi, Tokyo 191-0065 Japan
第二共同発明者	2 - 00	Full name of second joint inventor, if any	Tomohiro OHTA
第二共同発明者の署名	日付	Second inventor's signature	Date May 16, 2001
住所		Residence	TOKYO, JAPAN JPX
国籍		Citizenship	Japanese
私書箱		Post Office Address	c/o Teijin Limited, 1-1, Uchisaiwaicho 2-chome, Chiyoda-ku, Tokyo 100-0011 Japan

(第三以降の共同発明者についても同様に記載し、署名をする
こと) (Supply similar information and signature for third and
subsequent joint inventors.)

Japanese Language Declaration

第三共同発明者氏名 3-00	Full name of third joint inventor Keiji KOMORIYA
同発明者の署名 日付	Inventor's signature Keiji Komoriya Date May 18, 2001
住所	Residence TOKYO, JAPAN JPX
国籍	Citizenship Japanese
郵便の宛先	Post Office Address c/o Teijin Limited 1-1, Uchisaiwaicho 2-chome, Chiyoda-ku, Tokyo 100-0011 Japan
第四共同発明者 4-00	Full name of fourth joint inventor, if any Akira KUDO
同発明者の署名 日付	Inventor's signature Akira Kudo Date May 12, 2001
住所	Residence TOKYO, JAPAN JPX
国籍	Citizenship Japanese
郵便の宛先	Post Office Address 1-15-20, Nishi, Kunitachi-shi, Tokyo 186-0005 Japan

第五共同発明者氏名 5-00	Full name of fifth joint inventor Sunao TAKESHITA
同発明者の署名 日付	Inventor's signature Sunao Takeshita Date May 12, 2001
住所	Residence KANAGAWA, JAPAN JPX
国籍	Citizenship Japanese
郵便の宛先	Post Office Address 2-904, Lions Mansion Tsukimino, 2129-3, Shimotsuruma, Yamato-shi, Kanagawa 242-0001 Japan
第六共同発明者 6-00	Full name of sixth joint inventor, if any Keisuke KAJI
同発明者の署名 日付	Inventor's signature Keisuke Kaji Date May 12, 2001
住所	Residence KANAGAWA, JAPAN JPX
国籍	Citizenship Japanese
郵便の宛先	Post Office Address 108, Hodogaya Bldg., 4-9-6, Nagatsuta, Midori-ku, Yokohama-shi, Kanagawa 226-0027 Japan